Lab Dept:	Chemistry	
Test Name:	ALDOLASE	
General Information		
Lab Order Codes:	ALD	
Synonyms:	Fructose-biphosphate aldolase	
CPT Codes:	82085 - Aldolase	
Test Includes:	Aldolase level reported in U/L.	
Logistics		
Test Indications:	Useful for the detecting muscle diseases, such as Duchenne muscula dystrophy, dermatomyositis, polymyositis, and limb-girdle dystrophy.	
Lab Testing Sections:	Chemistry - Sendouts	
Referred to:	Mayo Medical Laboratories (Test: ALS)	
Phone Numbers:	MIN Lab: 612-813-6280	
	STP Lab: 651-220-6550	
Test Availability:	Daily, 24 hours	
Turnaround Time:	1 – 3 days, test set up Monday - Saturday	
Special Instructions:	Red No Gel tube required.	
Specimen		
Specimen Type:	Blood	
Container:	Red top (no gel) tube	
Draw Volume:	3 mL (Minimum: 1.5 mL) blood	
Processed Volume:	1 mL (Minimum: 0.5 mL) serum	
Collection:	Routine venipuncture	

Special Processing:	Lab Staff: Centrifuge specimen, remove serum aliquot into a screw- capped round bottom plastic vial. Hemolyzed specimens are not acceptable. Store and ship at refrigerated temperatures. Forward promptly.		
Patient Preparation:	None		
Sample Rejection:	Specimens other than serum; gross hemolysis; grossly icteric; gross lipemia; mislabeled or unlabeled specimens; specimens collected in gel tubes		
Interpretive			
Reference Range:	0 – 16 years:	<14.5 U/L	
	≥17 years:	<7.7 U/L	

-

Critical Values:N/ALimitations:N/AMethodology:Ultraviolet, KineticReferences:Mayo Medical Laboratories Web Page December 2017Updates:6/13/2007: Reference range update from MML. Previous ranges listed<br/>as 0-2 yrs: <16.3 U/L, 3-16 yrs: <8.3 U/L, >or= 17 yrs: <7.4 U/L.<br/>5/4/2009: Serum gel tubes are not acceptable for specimen collection.<br/>7/15/2013: Removed "fasting patient preferred".