Lab Dept: Serology

Test Name: CENTROMERE ANTIBODIES, IGG

**General Information** 

Lab Order Codes: CENA

**Synonyms:** Anti-Centromere Antibodies; ACA; CREST

**CPT Codes:** 83516 – Immunoassay for analyte other than infectious agent antibody

or infectious agent antigen; quaLitative or semi-quantitative, multiple

step method

**Test Includes:** Centromere antibody reported as negative or positive.

Logistics

**Test Indications:** Evaluating patients with clinical signs and symptoms compatible with

systemic sclerosis including skin involvement, Raynaud's phenomenon,

and arthralgias.

**Lab Testing Sections:** Serology - Sendouts

Referred to: Mayo Medical Laboratories (MML Test: CMA)

Phone Numbers: MIN Lab: 612-813-6280

STP Lab: 651-220-6550

**Test Availability:** Daily, 24 hours

**Turnaround Time:** 1-3 days, test set up Monday through Saturday

**Special Instructions:** N/A

Specimen

Specimen Type: Blood

**Container:** SST (Gold, marble or red)

**Draw Volume:** 1.5 mL (Minumum: 1.1 mL) blood

**Processed Volume:** 0.5 mL (Minimum: 0.35 mL) serum

**Collection:** Routine blood collection

**Special Processing:** Lab Staff: Centrifuge specimen, remove serum aliquot into screw-

capped round bottom plastic vial. Store and ship refrigerated. Forward

promptly.

Patient Preparation: None

Sample Rejection: Specimens other than serum, warm specimens, gross hemolysis, gross

lipemia, mislabeled or unlabeled specimens

## Interpretive

## **Reference Range:**

All Ages:	<1.0 U	Negative
	≥1.0 U	Positive

**Interpretation:** In various reported clinical studies, Anti-Centromere Antibodies occur in 50-96% of patients with CREST syndrome.

A positive test for Centromere Antibodies is strongly associated with CREST syndrome. The presence of detectable levels of centromere antibodies may antedate the appearance of diagnostic clinical features of CREST syndrome, in some cases by several years.

Critical Values: N/A

**Limitations:** Centromere antibodies have also been described in some patients with

primary biliary cirrhosis, and may occur in patients with rheumatoid

arthritis or lupus erythematosus.

Methodology: Multiplex Flow Immunoassay

References: Mayo Medical Laboratory Web Page January 2018

**Updates:** 1/6/2006: CPT 2006 changes

11/1/2007: Addition of units and numeric reporting along with negative and positive results. Change in method, previously listed as EIA. CPT

update, previously listed as 83516.

4/11/2011: CPT change from 88184 to 83520. 1/28/2016: CPT change from 83520 to 83516 1/17/2018: Collection container update.