Lab Dept: Serology

Test Name: VZV IgM ANTIBODY

General Information

Lab Order Codes: VZIGM

Synonyms: Anti-VZV Antibody IgM; VZV IgM Serology; Varicella-Zoster IgM Antibody

CPT Codes: 86787 – Antibody, varicella zoster (IgM)

Test Includes: Anti-VZV Antibody IgM reported as positive or negative.

Logistics

Test Indications: Intended for the qualitative detection of IgM antibodies to Varicella-Zoster

diagnosing acute-phase infection.

Lab Testing Sections: Immunology - Sendouts

Referred to: Mayo Medical Laboratories (MML Test: VZM)

Phone Numbers: MIN Lab: 612-813-6280

STP Lab: 651-220-6550

Test Availability: Daily, 24 hours

Turnaround Time: 1 - 3 days; test performed daily

Special Instructions: N/A

Specimen

Specimen Type: Blood

Container: SST (Gold, marble, or red)

Draw Volume: 1.5 mL (Minimum: 0.6 mL) blood

Processed Volume: 0.5 mL (Minimum: 0.2 mL) serum

Collection: Routine blood collection

Special Processing: Lab Staff: Centrifuge specimen, remove serum aliquot into a screw-topped

plastic vial. Store and ship at refrigerated temperatures. Forward promptly.

Patient Preparation: None

Sample Rejection: Hemolyzed specimens; lipemic specimens; mislabeled or unlabeled

specimens, specimens other than serum

Interpretive

Reference Range: Negative

A positive IgM result indicates a recent infection with VZV.

Critical Values: N/A

Limitations: A negative result does not rule out the diagnosis of VZV infection. Samples

collected early in the course of a VZV infection may not have detectable levels of antibody. In such cases, it is recommended that a second serum

sample be obtained 2-3 weeks later.

The performance characteristics with individuals vaccinated with VZV (OKA

Strain) have not been established.

The test must be performed on serum. The use of whole blood, plasma or

cord blood has not been established.

Positive results from cord blood or neonates should be interpreted with

caution.

Results from immunocompromised patients should be interpreted with

caution.

Methodology: Immunofluorescence Assay (IFA)

References: <u>Mayo Medical Laboratories</u> January 2018

Updates: 9/12/2005: Testing turnaround time previously listed as 1-6 days, testing

performed 1 time per week.

2/1/2006: Test moved from Children's Hospitals and Clinics of Minnesota

Laboratories to Mayo Medical Laboratories. 1/17/2018: Collection container update.