## Lab Dept: Microbiology

| Test Name:            | LEGIONELLA CULTURE  |
|-----------------------|---|
| General Information   |   |
| Lab Order Codes:      | LEGC  |
| Synonyms:             | Culture, Legionella pneumophila   |
| CPT Codes:            | 87081 – Culture, presumptive, pathogenic organisms, screening only 87077 – Aerobic isolate, additional methods required for definitive identification of isolates |
| Test Includes:        | Culture for Legionella pneumophila.   |
| Logistics             |   |
| Lab Testing Sections: | Microbiology  |
| Referred to:          | Minnesota Department of Health  |
| Phone Numbers:        | MIN Lab: 612-813-5866   |
|                       | STP Lab: 651-220-6555   |
| Test Availability:    | Daily, 24 hours   |
| Turnaround Time:      | Positive results are usually generated between 2 - 5 days. Negative cultures are final at 14 days.  |
| Special Instructions: | Specimen site and date/time of collection are required for processing.  |
| Specimen              |   |
| Specimen Type:        | Bronchoscopy, blood, bone marrow, lung aspirate, lung biopsy, pericardial fluid, peritoneal fluid, pleural fluid, tracheal aspirates or sputum                    |
| Container:            | Sterile container   |
| Volume:               | 1 mL fluid  |
| Collection:           | Bronchoscopy:   |
|                       | <b>1.</b> Specimen obtained by physician through the biopsy channel of the bronchoscope.  |

- 2. Transfer specimen into a luki tube.
- **3.** Transport to laboratory immediately at room temperature.

## Lung Aspirates, Pleural Fluid:

- **1.** Disinfect overlying skin with 2% tincture of iodine.
- **2.** Obtain specimen via percutaneous needle aspiration or surgery.
- 3. Place in sterile container.
- **4.** Transport to the Laboratory immediately at room temperature.

## Lung Biopsy:

- 1. Any visible amount.
- 2. Submit in sterile container without formalin.
- 3. Lab Staff: Add a few drops of sterile saline to prevent drying.

## Sputum (Expectorate):

|                     | <ol> <li>Collect early morning specimen under the direct supervision<br/>of a nurse or a physician.</li> <li>Have patient rinse or gargle with water to remove superficial<br/>flora.</li> <li>Instruct patient to cough deeply to produce a lower<br/>respiratory specimen.</li> <li>Do not submit saliva.</li> </ol>  |
|---------------------|---|
|                     | Sputum (Induced):   |
|                     | <ol> <li>Have patient rinse mouth with water after brushing gums and tongue.</li> <li>With the aid of a nebulizer, have patients inhale ~25 mL of 3 to 10% sterile saline.</li> <li>Collect the induced sputum in a sterile container.</li> </ol>   |
| Transport/Storage:  | Transport to the Microbiology Laboratory immediately at room temperature.   |
| Special Processing: | Refrigerate specimens that cannot be processed within 30 min. If processing is delayed more than 24 hours, freeze specimen at -70°C.  |
| Sample Rejection:   | Throat and nasopharyngeal specimens are not acceptable due to the<br>presence of normal respiratory flora. Specimen with a transit time<br>exceeding 1 hour after collection; specimen not submitted in<br>appropriate transport container; improperly labeled specimen;<br>insufficient volume; external contamination. If an unacceptable<br>specimen is received, the physician or nursing station will be notified<br>and another specimen will be requested before the specimen is<br>discarded. |
| Interpretive        |   |

**Reference Range:** 

No Legionella pneumophila isolated.

| Alert Values:           | The physician will be notified of positive results.  |
|-------------------------|--|
| Limitations:            | Sputum (expectorated), tracheal aspirates, and other specimens having<br>normal respiratory flora are subject to bacterial overgrowth and may<br>mask growth of Legionella. Sensitivity of cultures is relatively low (50%<br>to 80%), however, specificity approaches 100%.   |
| Methodology:            | Culture  |
| Additional Information: | The family, <i>Legionellaceae</i> , are ubiquitous, gram-negative, motile, fastidious, aerobic bacilli. Disease outbreaks have been associated with exposure of susceptible individuals to water sources in which legionellae have grown. <i>Legionella</i> sp. cause respiratory illness manifested primarily by pneumonia or Pontiac fever, a non-pneumonic, influenza-like illness. During an American Legion Convention in Philadelphia in 1976, an epidemic of pneumonia caused 34 deaths. Sputum characterized by acute inflammatory features, without a classical pattern of bacteria, may represent <i>Legionella</i> , influenza, or respiratory syncytial virus. |
| References:             | <ul> <li>Cook, JH, and M Pezzlo (1992). Specimen receipt and accessioning.</li> <li>Section 1. Aerobic bacteriology, 1.2.1-4. In HD Isenberg (ed) Clinical Microbiology Procedures Handbook. American Society for Microbiology, Washington DC</li> <li>Miller, J Michael (1999) A Guide To Specimen Management in Clinical Microbiology, American Society for Microbiology, Washington DC</li> <li>Miller, J Michael, and HT Holmes (1999) Specimen Collection, Transport, and Storage In PR Murray et al, (ed), Manual of Clinical Microbiology, 7<sup>th</sup> edition, American Society for Microbiology, Washington DC, pp 33-104</li> </ul>                           |
| Test Updates:           | 6/2/2014: DFA no longer performed as part of this test.  |