
Lab Dept: Urine/Stool

Test Name: 17-HYDROXYPROGESTERONE, URINE

General Information

Lab Order Codes: 17HU

Synonyms: 17-Hydroxy Progesterone, Urine; 17-OH Progesterone, Urine

CPT Codes: 83498 – Hydroxyprogestone, 17-d

Test Includes: Urine 17-Hydroxy Progesterone is measured by radioimmunoassay following extraction of specimens.

Logistics

Test Indications: 17-Hydroxy Progesterone is a steroid derived primarily from enzymatic metabolism of Progesterone and 17-Hydroxy Pregnenolone. It is converted enzymatically to Androstenedione and 11-Deoxycortisol. It is produced in both the gonads and adrenal glands. It is excreted into the urine in conjugated and unconjugated forms of 17-Hydroxy Progesterone and as Pregnanetriol. This assay measures the total of the conjugated and unconjugated forms. It is stimulated by ACTH and suppressed by Dexamethasone. Levels of urine 17-Hydroxy Progesterone are greatly increased in patients with Polycystic Ovarian Disease and Congenital Adrenal Hyperplasia and show exaggerated responses to ACTH in these cases. 17-Hydroxy Progesterone is the marker steroid for determining cases of 21a-Hydroxylase Deficient Congenital Adrenal Hyperplasia. Urine levels are frequently elevated in patients with idiopathic hirsutism.

Lab Testing Sections: Urine/Stool - Sendouts

Referred to: Mayo Medical Laboratories (MML Test: ZW71 - Misc) and forwarded to Inter Science Institute

Phone Numbers: MIN Lab: 612-813-6280

STP Lab: 651-220-6550

Test Availability: Daily, 24 hours

Turnaround Time: 5 – 9 days; performed Monday - Friday

Special Instructions: Refrigerate or keep specimen on ice during 24 hour collection. There is no preservative. [See Patient Preparation.](#)

Specimen

Specimen Type:	Urine, 24 hour collection
Container:	Plastic, leak-proof urine collection container
Draw Volume:	Entire collection
Processed Volume:	10 mL (Minimum: 5 mL) urine from a well-mixed 24 hour collection
Collection:	Routine 24 hour urine collection, refrigerate during collection
Special Processing:	Lab Staff: Mix 24 hour specimen well. Measure and record total urine volume. Remove aliquot into a screw-capped round bottom plastic vial. Provide total 24 hour volume. Store and ship at frozen temperatures. Forward promptly.
Patient Preparation:	Patient should not be on any Corticosteroid, ACTH, steroid, estrogen, or gonadotropin medication, if possible, for at least 48 hours prior to start of urine collection.
Sample Rejection:	Specimens other than urine, mislabeled specimens or unlabeled specimens

Interpretive

Reference Range:	17-Hydroxyprogesterone:	
	Age	Range
	Adult Males:	Up to 2.0 ug/24 hours
	Adult Females:	Up to 4.5 ug/24 hours

Critical Values:	N/A
Limitations:	N/A
Methodology:	Radioimmunoassay (RIA) following extraction of specimens
References:	Mayo Medical Laboratory Web Page November 2017 Inter Science Institute November 2017
Updates:	11/13/2017: Updated reference range and order codes.