


Aim: To minimize use of unnecessary pre-medications for transfusion reactions and to clarify those patients that require pre-medications.



EXCLUSION GUIDELINES
Patients **excluded** from this guideline:
Patients with history of:

- Hemolytic transfusion reaction
- Transfusion Related Acute Lung Injury (TRALI)
- Delayed hemolytic transfusion reaction
- Transfusion Associated Circulatory Overload (TACO)
- Septic transfusion reaction

If questions or concerns, please contact the on-call hematologist via Amion.

NOTE 1:
As documented in chart, problem-list, or in discussion with patient and caregivers

NOTE 2:
Mild symptoms isolated to fever, chills/rigors without evidence of hemolysis

NOTE 3:
Allergic Reaction Definitions

- **Mild:** urticarial, pruritis, and/or localized angioedema within 4 hours of transfusion
- **Severe:** respiratory compromise (tachypnea, wheezing, dyspnea) and/or hypotension

Aim: To minimize use of unnecessary pre-medications for transfusion reactions and to clarify those patients that require pre-medications.

REFERENCES:

1. Geiger TL, Howard SC. Acetaminophen and diphenhydramine premedication for allergic and febrile nonhemolytic transfusion reactions: good prophylaxis or bad practice? *Transfus Med Rev.* 2007 Jan;21(1):1-12. doi: 10.1016/j.tmr.2006.09.001. PMID: 17174216; PMCID: PMC1868404.
2. Goel R, Tobian AAR, Shaz BH. Noninfectious transfusion-associated adverse events and their mitigation strategies. *Blood.* 2019 Apr 25;133(17):1831-1839. doi: 10.1182/blood-2018-10-833988. Epub 2019 Feb 26. PMID: 30808635.
3. Martí-Carvajal AJ, Solà I, González LE, Leon de Gonzalez G, Rodriguez-Malagon N. Pharmacological interventions for the prevention of allergic and febrile non-haemolytic transfusion reactions. *Cochrane Database Syst Rev.* 2010;2010(6):CD007539. Published 2010 Jun 16. doi:10.1002/14651858.CD007539.pub2
4. McCormick M, Triulzi D. The use of premedications for platelet transfusions in pediatric patients. *Hematology Am Soc Hematol Educ Program.* 2020 Dec 4;2020(1):523-526. doi: 10.1182/hematology.2020000165. PMID: 33275693; PMCID: PMC7727506.
5. Sanders RP, Maddirala SD, Geiger TL, Pounds S, Sandlund JT, Ribeiro RC, Pui CH, Howard SC. Premedication with acetaminophen or diphenhydramine for transfusion with leucoreduced blood products in children. *Br J Haematol.* 2005 Sep;130(5):781-7. doi: 10.1111/j.1365-2141.2005.05670.x. PMID: 16115137.
6. Tobian AA, King KE, Ness PM. Transfusion premedications: a growing practice not based on evidence. *Transfusion.* 2007 Jun;47(6):1089-96. doi: 10.1111/j.1537-2995.2007.01242.x. PMID: 17524101.