
Lab Dept: Chemistry

Test Name: GROWTH HORMONE, SINGLE, SERUM

General Information

Lab Order Codes: HGH

Synonyms: HGH; Somatotropin

CPT Codes: 83003 - Growth hormone, human (HGH)

Test Includes: Single growth hormone level reported in ng/mL.

Logistics

Test Indications: Excess Growth Hormone secretion is associated with acromegaly in adults or pituitary gigantism in children, while a deficiency may cause short stature in children.

Lab Testing Sections: Chemistry – (performed on Minneapolis Campus)

Phone Numbers: MIN Lab: 612-813-6280

STP Lab: 651-220-6550

Test Availability: Daily, 24 hours

Turnaround Time: Performed daily

Special Instructions: Follow physician's protocol for administering provocative stimuli and drawing specimens, if appropriate. See [Patient Preparation](#).

Specimen

Specimen Type: Blood

Container: Red NO GEL, preferred
Alternate: SST (gold or marble)

Draw Volume: 2.1 mL (Minimum: 1.0 mL) blood

Processed Volume: 0.7 mL (Minimum: 0.3 mL) serum

Collection: Routine blood collection

Special Processing: Lab Staff: Centrifuge specimen within 1 hour of draw, remove serum aliquot into a screw-capped round bottom plastic vial. Freeze immediately. Ship to Minneapolis lab frozen.

Patient Preparation: See [Special Instructions](#). In addition, patient must be fasting and at complete rest 30 minutes before blood collection.

Sample Rejection: Mislabeled or unlabeled specimens; improper storage; gross hemolysis; incorrect sample type

Interpretive

Reference Range:	Age	Result Range
	0 - 1 day	5 - 53 ng/mL
	1 day – 13 days	5 - 27 ng/mL
	14 days - 12 months	2 - 10 ng/mL
	>12 months:	0 – 6 ng/mL
	Response Testing (Children and Adults):	Following provocative stimuli, a normal response is a rise to 10 ng/mL or greater. Intermediate values between 7 ng/mL and 10 ng/mL may indicate a partial deficiency and require further evaluation.

Critical Values: N/A

Limitations: Patients on hGH therapy may develop human antibodies to hGH that will interfere with the test and cause falsely low levels.

This method is sensitive to biotin interference. Unexpected results that do not match clinical presentation should be repeated at least 24 hours after the last dose of supplemental biotin, which includes some multivitamins.

Methodology: Chemiluminescent competitive immunoassay

References: IDS iSYS HGH Reagent Package Insert. Revised January 31, 2017, IS-3700 PL V04

National Committee for Clinical Laboratory Standards (NCCLS) (2002) Clinical Laboratory Technical Procedure Manuals; Approved Guideline-Fourth Edition GP2-A4

Expected Values and S.I. Unit Conversion Tables, 2nd Edition,
Endocrine Sciences, 4301 Lost Hills Rd, Calabasas Hills, CA 91301

Biorad Lyphochek Immunoassay Plus Control Product Insert

Burtis, CA, Ashwood, ER, Bruns, DE, ed (2006) Tietz Textbook of
Clinical Chemistry and Molecular Diagnostics. 4th Edition, Philadelphia:
W.B. Saunders

Jacobs and DeMott Laboratory Test handbook (2001) 5th Edition, Lexi-
comp Inc

Updates:

10/7/2016: Updated tube type.

6/26/2017: Added alternate tube types.

6/16/2018: New method ISD iSYS– inhouse test, sample type change