

---

**Lab Dept:** Microbiology/Virology

**Test Name:** MYCOPLASMA HOMINIS PCR

---

**General Information**

**Lab Order Codes:** MYHP

**Synonyms:** PCR, *Mycoplasma hominis*

**CPT Codes:** 87798 – Infectious agent detection by nucleic acid (DNA or RNA), amplified probe technique

**Test Includes:** Rapid, sensitive PCR analysis of submitted specimen for detection of *Mycoplasma hominis* and reported as positive or negative.

---

**Logistics**

**Lab Testing Sections:** Microbiology - Sendouts

**Referred to:** Mayo Medical Laboratories (Test: MHRP)

**Phone Numbers:** MIN Lab: 612-813-6280

STP Lab: 651-220-6550

**Test Availability:** Daily, 24 hours

**Turnaround Time:** 3 - 4 days

**Special Instructions:** **Specimen site** and **date/time of collection** are required for specimen processing.

---

**Specimen**

**Specimen Type:** **Swabs:** Cervix, Urethra or Vaginal; upper respiratory sources (only infants <3 months only: nasopharynx, nose, throat)

**Fluids:** Reproductive fluids, Respiratory specimens, pleural fluid, pericardial fluid, lung or synovial fluid

**Urine, Kidney stones, Ureter, Tissue** (placenta, Products of conception, Genitourinary, Respiratory, Wound)

**Plasma:** EDTA derived

**Whole blood:** EDTA

**Container:**

**Swabs:** Swab transport **Note:** Wooden shaft, cotton swabs are not acceptable.

**Fluids/Urine/Kidney stones/Tissue:** Sterile container

**Plasma:** EDTA tube

**Whole blood:** EDTA tube

**Draw Volume:**

**Swabs:** 1 swab

**Fluids:** 2 mL (Minimum: 1 mL) fluid

**Urine:** 10 mL (Minimum: 2 mL) urine

**Kidney stones:** Entire specimen

**Tissue:** 5mm fresh tissue

**Plasma:** 3 mL blood

**Whole blood:** 1 mL blood in EDTA tube

**Collection:**

**Do not** collect specimens on wooden shaft swabs because the shaft is toxic to these organisms.

**Swab specimens: Do not** collect specimens on wooden shaft swabs because the shaft is toxic to these organisms.

**Cervical, Vagina, or Urethral Specimens:**

1. Obtain specimen from infected site on swab.
2. Collect vaginal or throat specimen by swabbing back and forth over the mucosa to maximize recovery of cells. Collect urethra and cervical specimen by inserting swab 1 cm to 3 cm and rotating 360 degrees.
3. Place the swab back into the swab cylinder and indicate specimen source on label.
4. Deliver to Laboratory immediately.

Lab Staff:

1. Write the specimen source on the label.
2. Send specimen refrigerated. Maintain sterility and forward promptly.

**Amniotic Fluid, Prostatic Secretions, Respiratory Specimens (<3 months old), Semen, Reproductive Drainage/fluid or Synovial Fluid:**

1. Collect 2 mL specimen in a sterile container and deliver to laboratory immediately. Include specimen source.

Lab Staff:

1. Write the specimen source on the label.
2. Send specimen refrigerated. Maintain sterility and forward promptly.

**Urine: Clean catch, Mid-stream specimen:**

**Males:**

1. Clean glans with soap and water.
2. Rinse area with wet gauze pads.
3. While holding foreskin retracted, collect the first 2-10 mL from urine stream in a sterile container.
4. Send specimen to lab.

## Lab Staff:

1. Maintain sterility, refrigerate specimen and forward promptly.

**Females:**

1. Thoroughly clean urethral area with soap and water.
2. Rinse area with wet gauze pads.
3. While holding labia apart, collect the first 2-10 mL from urine stream in a sterile container.
4. Send specimen to lab.

## Lab Staff:

1. Maintain sterility, refrigerate specimen and forward promptly.

**Kidney stones:** Stone collection

## Lab Staff:

1. Maintain sterility, refrigerate specimen and forward promptly

**Tissue:** Collection by physician or medical provider

## Lab Staff:

1. Maintain sterility, refrigerate specimen and forward promptly

**Plasma:** Routine venipuncture

## Lab Staff:

1. Centrifuge specimen, maintain sterility and transfer a 1 mL aliquot to a sterile tube, refrigerate specimen and forward promptly.

**Whole blood:** Routine venipuncture

## Lab Staff:

1. DO NOT centrifuge, specimen should remain in original collection container, refrigerate specimen and forward promptly.

**Transport/Storage:** Transport all specimens to the lab at refrigerated temperatures.

**Lab Staff:**

**Swabs:** Swab in original swab cylinder.

**Fluid specimens (other than urine):** Place 1-2 mL fluid into a sterile container.

**Urine:** Send undiluted urine in sterile container

**Kidney stone:** Submit in sterile container

**Tissue:** Submit in sterile container.

All specimens should be **refrigerated** for transport to reference lab.

**Note:** Swabs and Fluids stored in M5 transport media will be accepted and tested, but the preferred specimen is the direct specimen without transport media.

**Sample Rejection:** Specimen not submitted in appropriate transport container; improperly labeled specimen; insufficient volume; external contamination; warm specimens. If an unacceptable specimen is received, the physician or nursing station will be notified and another specimen will be requested before the specimen is discarded.

---

***Interpretive***

**Reference Range:** Negative (reported as positive or negative)

**Limitations:** Interfering substances may affect the accuracy of this assay; results should always be interpreted in conjunction with clinical and epidemiological findings.

This test does not detect other mycoplasmas or ureaplasmas (including *Mycoplasma pneumoniae*, a common cause of community acquired pneumonia).

**Methodology:** Real-Time Polymerase Chain Reaction (PCR) using LightCycler and Fluorescent Resonance Energy Transfer (FRET)

**Additional Info:** *Mycoplasma hominis* causes genitourinary tract diseases. Importantly, infection of infants with *M. hominis* occurs during passage through an infected birth canal. Infection may result in neonatal meningitis and pneumonia. In adults, *M. hominis* is a recognized cause of acute pelvic inflammatory disease, postpartum fever and pyelonephritis.

**References:** [Mayo Medical Laboratories](#) September 2015

**Updates:** 8/13/2015: Added kidney stones and fresh tissue  
9/29/2015: Added Plasma and Whole blood specimens

