Lab Dept:

# Microbiology

## Test Name: OVA AND PARASITE EXAMINATION, STOOL

| <b>General Information</b> | ormation |
|----------------------------|----------|
|----------------------------|----------|

| Lab Order Codes:                          | OAP  |
|---|--|
| Synonyms:                                 | O & P Exam, stool; Stool for Ova and Parasites; Feces for Ova and Parasites; Parasitology examination  |
| CPT Codes:                                | 87177 – Ova and parasites, direct smears, concentration and identification<br>87209 – Smear, primary source with interpretation; complex special stain for ova and parasites |
| Test Includes:                            | Examination of stool for intestinal parasites by direct/concentrated microscopic exam and trichrome stain.   |
|   | This test <b>does not</b> include: <i>Cryptosporidium</i> , <i>Cyclospora</i> , <i>Microsporidium</i> or Pinworm.  |
|   | If only <i>Cryptosporidium</i> or <i>Giardia</i> are requested, refer to specific listing for <u>Cryptosporidium/Giardia FA.</u>   |
|   |  |
|   | If pinworm is requested, refer to Pinworm Preparation.   |
| Logistics                                 | If pinworm is requested, refer to Pinworm Preparation.   |
| <i>Logistics</i><br>Lab Testing Sections: | If pinworm is requested, refer to <u>Pinworm Preparation.</u>  |
| -   |  |
| Lab Testing Sections:                     | Microbiology   |
| Lab Testing Sections:                     | Microbiology<br>MIN Lab: 612-813-5866  |
| Lab Testing Sections:<br>Phone Numbers:   | Microbiology<br>MIN Lab: 612-813-5866<br>STP Lab: 651-220-6555   |

| Special Instructions: | <ul> <li>Specimen site and date/time of collection are required for specimen processing.</li> <li>Provide clinical diagnosis and travel history or suspected parasite on request form. If pinworm is requested, refer to <u>Pinworm Preparation.</u></li> </ul>  |
|-----------------------|--|
|                       | <b>Warning</b> : Any stool collected from a patient infected with parasites is highly infectious. Use extreme caution when <i>Entamoeba histolytica</i> , <i>Hymenolepis nana</i> , and <i>Taenia</i> sp. are suspected.   |
| Specimen              |  |
| Specimen Type:        | Fresh random stool or aspirated stool; early morning sputum; mid-day urine   |
| Container:            | Onsite collections: Plastic, leak-proof container  |
|                       | <b>Offsite collections:</b> O&P preservative kit (available from Materials, Storeroom Item# 14574)   |
|                       |  |
| Volume:               | 5 - 10 mL liquid stool or 5 – 10 grams of formed stool   |
|                       | 100 mL urine – entire mid-day urine collection with last voided portion containing blood and mucous  |
|                       | 1-3 mL sputum  |
|                       | Note: If O&P kit is used, make sure to add specimen to "fill" lines.   |
| Collection:           | Recommendations:   |
|                       | <ul> <li>Maximum of 3 specimens, collected on separate days within a 10-day time period.</li> <li>Do not submit more than one specimen per day. Many organisms do not appear in stool specimens in consistent numbers on a daily basis; few or no parasites may be passed on one day, with many passed the next day.</li> <li>Parasite exams on patients hospitalized &gt;3 days are not productive and should not be ordered unless special circumstances exist.</li> </ul> |
|                       | Fresh Stool:   |
|                       | Onsite collections ONLY:<br>1. Collect stool in a clean, dry bedpan or on a newspaper over the   |

toilet. Do not contaminate specimen with urine, residual soap or disinfectants, which will destroy amoebae.

2. Transfer to a plastic, leak-proof container.

**3**. If a delay >1 hour in transport is expected, place specimen in O&P preservative kit. <u>Refer to Special Processing</u>.

**4.** Those portions of stool containing blood and mucous are especially significant and should be transferred into the container.

**5.** Specimens in diapers are **not** acceptable. Pediatric patients with severe diarrhea may use a U bag collection system. Place the bag over the anal area in an attempt to retrieve the specimen before it soaks into the diaper. The diaper can also be reversed with the plastic side toward the skin to prevent the specimen from soaking into the diaper. Transfer specimen into a plastic, leak-proof container.

#### Offsite collections:

See Special Processing for O&P preservation

#### Sputum (Expectorate):

#### **Onsite collections ONLY:**

**1.** Collect early morning specimen under the direct supervision of a nurse or physician.

2. Have patient rinse or gargle with water to remove superficial flora.

**3.** Instruct patient to cough deeply to produce a lower respiratory specimen.

**4.** Exam specimen to make sure it contains thick mucus. Do not submit saliva.

**5.** Transport directly to the laboratory (< or =15 min) since specimens must be examined with one hour of collection.

### Urine:

#### **Onsite collections ONLY:**

**1.** Collect mid-day urine specimen in a sterile container. Peak egg excretion occurs between noon and 3 pm.

**2.** For patients with hematuria, eggs are associated with the terminal (last voided) portion of the specimen containing blood and mucus.

**3.** Transport directly to the laboratory (< or =15 minutes) since specimens must be examined within one hour of collection.

| Special Processing:  | Instructions for O&P preservative kit: 2 vials, one containing 10% formalin (pink) and one containing Zn-PVA (gray).  |
|----------------------|---|
|                      | Caution: The liquid preservative in the vials is poisonous.   |
|                      | <ol> <li>Fill each vial by using the spoon built into the lid of the vial and<br/>transferring small scoopfuls of stool from areas which appear bloody,<br/>slimy or watery until the contents rise to the "Fill Here" red line. Do not<br/>overfill.</li> <li>If the stool is formed, sample small amounts from each end, sides<br/>and the middle.</li> </ol>   |
|                      | <b>3.</b> Mix the contents of the vials with the spoon. Screw cap on tightly and shake the vial vigorously until the contents are well mixed. Make sure there is no leakage.  |
|                      | <ul> <li>4. Label vials with patient's name, date and time of collection.</li> <li>5. Store vials at room temperature.</li> <li>6. Return collection kit to laboratory within 72 hours.</li> </ul>  |
| Storage /Transports  |   |
| Storage/Transport:   | <b>Onsite collections:</b> Transport to the laboratory immediately.   |
|                      | <b>Offsite collections: Do not refrigerate</b> , store at room temperature.<br>Specimens must be promptly transported to the laboratory, with the next available courier.   |
| Patient Preparation: | <b>Do not</b> use antacids, barium, bismuth, antidiarrheal medication, antibiotics, or oily laxatives for 5-10 days prior to collection.  |
| Sample Rejection:    | <b>No diapers accepted</b> . Unpreserved specimens (fresh) will be rejected with a transit time exceeding 1 hour after collection; specimens containing interfering substances (e.g., antidiarrheal compounds, antibiotics, antacids, oils, bismuth, or barium); more than one specimen per day; specimen not submitted in appropriate transport container; improperly labeled specimen; insufficient volume; overfilled or underfilled O&P kits; specimen contaminated with toilet water or urine; external contamination. If an unacceptable specimen is received, the physician or nursing station will be notified and another specimen will be requested before the specimen is discarded. |
| Interpretive         |   |
| Reference Range:     | No ova or parasites seen  |
| Limitations:         | One negative result does not rule out the possibility of parasitic infestation.   |
| Methodology:         | Concentrated microscopic exam and trichrome stain   |

| Additional Information: | Amebae and other parasites cannot be seen in stool containing barium.<br>Formed stool are more likely to contain amebic cysts and helminth<br>eggs. Soft or liquid stools are more likely to contain trophozoites of the<br>protozoa. Liquid stools must be examined within 30 minutes of<br>passage, not 30 minutes from the time the specimen arrives in the<br>laboratory. Soft (semi-formed) specimens should be examined within 1<br>hour of passage. If these time frames are not possible, the specimen<br>should be preserved in 10% formalin and Zn-PVA. Formalin will<br>preserve the protozoan cysts, larvae and eggs of the helminthes. Zn-<br>PVA will preserve the trophozoite stage of the protozoa. Immediate<br>examination of formed stools is not as critical as the protozoan cysts will<br>remain intact generally up to 24 hours. |
|-------------------------|---|
| References:             | Cook, JH, and M Pezzlo (1992). Specimen receipt and accessioning.<br>Section 1. Aerobic bacteriology, 1.2.1-4. In HD Isenberg (ed) Clinical<br>Microbiology Procedures Handbook. American Society for Microbiology,<br>Washington DC  |
|                         | Miller, J Michael (1999) A Guide To Specimen Management in Clinical Microbiology, American Society for Microbiology, Washington DC  |
|                         | Miller, J Michael, and HT Holmes (1999) Specimen Collection,<br>Transport, and Storage In PR Murray et al, (ed), Manual of Clinical<br>Microbiology, 7 <sup>th</sup> edition, American Society for Microbiology,<br>Washington DC, pp 33-104  |
|                         | Garcia, LS, RY Shimizu and P Deplazes (2003) Specimen Collection,<br>Transport, and Processing; Parasitology In PR Murray et al, (ed),<br>Manual of Clinical Microbiology, 8 <sup>th</sup> edition, American Society for<br>Microbiology, Washington DC, p 1908   |
| Updates:                | <ul> <li>1/20/2009: Amended formed stool volume, previously listed as 1 gram.</li> <li>11/20/2014: Offsite information added.</li> <li>2/23/2017: CPT update, stain method previously listed s 88312.</li> <li>2/19/2020: Updated Storage/Transport.</li> </ul>   |