

IV Medicine at Home: IV Push Method

Patient Name : _____ **MRN** _____ **DOB** _____

Medication Information

Name of Medication: _____ Dose: _____

Medication stable at room temperature for: _____

Give medication every _____ hours

Times to give medication:

| | | | | | |
|--|--|--|--|--|--|
| | | | | | |
|--|--|--|--|--|--|

Normal saline flush solution

Amount to give: _____ ml

When to give it:

- before and after giving the medication
- _____

Heparin flush solution

Solution: _____ units/ml

Amount to give: _____ ml

When to give it:

- After giving the medication and normal saline flush
- _____

Preventing infection

It is very important to prevent infection. Failure to do so might require removal of the central venous catheter. The nurse will show you how to keep your supplies clean, so no germs will enter the catheter and cause an infection.

Cleanliness is a must!

Each time you handle supplies or perform a procedure, do the following steps:

1. Prepare a clean work space area by cleaning a solid surface with household cleanser.
2. Clean hands well, using one of the following methods:
 - Use an alcohol hand sanitizer according to directions.
 - If hands are dirty or alcohol hand sanitizer is not available, wash hands for *at least 20 seconds* with soap, rubbing all surfaces briskly, including fingernails. Use a paper towel or clean hand towel to dry your hands, and then use the towel to turn off the faucet.
3. Before *each time* you flush or attach a new syringe/tubing to the injection cap, scrub the injection cap for 30 seconds with a fresh chlorhexidine wipe, using friction. Let it dry. Do not fan or blow on it.

Getting ready

1. Make sure the medication is at room temperature
 - Remove the medication from the refrigerator as directed. Your homecare nurse will discuss with you how early this should be removed to make sure it is safe to use.
 - Do not use a microwave, oven, or water to warm.
2. Check the label on the syringe and verify:
 - Child's name
 - Medication/Dose
 - Expiration date

Connecting Medication/Fluids to Patient

Review of S.A.S.H. method:

1. Flush with normal **S**aline
2. **A**dminister medication or fluid
3. Flush with normal **S**aline
4. Flush with **H**eparin



Note: Every time you flush the IV line with normal saline or heparin, use the "**push-and-pause**" method: push a little solution, then pause for 1 to 2 seconds, then push a little more, pause, and so on. This method cleans the inside of the IV line. Your home care nurse will show you how to do this.

1. Gather supplies:

- chlorhexidine wipes
- normal saline syringes or dextrose solution syringes (2)
- heparin syringe
- IV medicine/fluids you will be giving
- non-sterile gloves
- disinfectant cap

2. Put supplies in order of use on the work surface.

3. Check the labels of the normal saline and heparin syringes, and any other medicines or fluids before giving them. Make sure they are the right medicines.

- A medicine is good until the end of the day on the day it expires. Throw it away sooner if you see sediment in it or if you question its cleanliness. If you do throw a medicine away, be sure to call the home care pharmacy for a replacement dose.

Preparing pre-filled syringes:

All pre-filled syringes have an air bubble in them (normal saline, heparin). Remove the tip cover from the syringe and point the syringe upward, pull back slightly on plunger and squirt air bubble out.

4. Put on non-sterile gloves.
5. Remove disinfectant cap **OR** if disinfectant cap not in place, scrub the injection cap with a chlorhexidine wipe for 30 seconds and let it dry.
6. Attach normal saline syringe tip into the injection cap.
7. Unclamp the catheter.
8. Flush with normal saline using the push-and-pause method. When syringe is empty, remove it from the injection cap.
9. If you are not giving medicines at this time, go to step 1 of the next section
10. If you are giving medicines, scrub injection cap with a new chlorhexidine wipe for 30 seconds and let it dry.
11. Connect the IV tubing to the patient's IV line.
12. Open all clamps.
13. Press the desired start button ("Normal", "Slow", or "Extra-Slow") to begin the infusion. A blinking green light shows that the pump is running.
14. Put the ESP pump on a nearby surface.

At end of infusion or when medication complete:

1. Disconnect the tubing from the injection cap.
2. Scrub the injection cap with a new chlorhexidine wipe for 30 seconds and let it dry.
3. Attach normal saline syringe or dextrose solution syringe and flush using the push-and-pause method. When syringe is empty, remove it from the injection cap.
4. Scrub the injection cap with a new chlorhexidine wipe 30 seconds and let it dry.
5. Attach heparin syringe and flush using the push-and-pause method. When syringe is empty, remove it from the injection cap.
6. Clamp the line.
7. Apply disinfectant cap to injection cap.

Questions?

If you have any questions, please call your home care nurse.

Reviewed by Home Care & Home Care Pharmacy 12/2016.